The Solid State

WS 2013/14

Lectures (Tuesday & Friday)

Paul van Loosdrecht
Optical Condensed Matter Physics, II. Physikalisches Institut
pvl@ph2.uni-koeln.de

Website II. Physikalisches Institut
www.ph2.uni-koeln.de
http://www.ph2.uni-koeln.de/527.html

Last time:

- Reciprocal lattice and diffraction cont'd
- Phonons

Today:

- Phonons cont'd
- Thermal properties

Phonons

Total lattice energy $U_{total} = \sum_{<ij>} U_{ij}(\vec{R}_j - \vec{R}_i)$

Stability: $\left.\frac{\partial U_{total}}{\partial R_j}\right|_{R_j = R_j^0} = 0 \Rightarrow$ Equilibrium coordinates

<u>Harmonic approximation:</u> $F_j = -\frac{\partial U_{total}}{\partial R_j}$

phonons

Classical: $u_s(t) = u_k(t)\, e^{i\, kas}$

EOM: $m\ddot{u}_k(t) = -C_k u_k(t)$ with $C_k = m\omega_k^{\,2}$

Quantum states of elastic waves:

$$\left\{ -\frac{\hbar}{2m}\frac{\partial^2}{\partial u_k^{\,2}} + \frac{1}{2} m\omega_k^{\,2} u_k^{\,2} \right\} \cdot \psi_k(u_k) = E_k \cdot \psi_k(u_k)$$

$$\psi_k(u_k, t) = e^{i(E_k/\hbar)t} \cdot \psi_k(u_k) \qquad E_k = (n + \frac{1}{2}) \cdot \hbar\omega_k$$



See appendix C: quantization of elastic waves: phonons

Three dimensions

3-dimensional crystal: 3 degrees of freedom per atom
s atoms per primitive cell: 3 acoustical branches
3*s*-3 optical branches

Optical modes

Acoustical modes



Longitudinal: u // k

Transverse: u ⊥ k

Properties of phonons

Any number can occupy the same vibrational mode u_k

Phonons are *bosons*

Thermal occupation given by Planck's distribution

$$\langle n_k \rangle = \frac{1}{e^{(\hbar\omega_k / k_b T)} - 1}$$

Zero-point energy: $E_0 = \sum_k \frac{1}{2}\hbar\omega_k$

Energy of 1 phonon: $\hbar\omega_k$

"Momentum" of 1 phonon: $\hbar k$

Phonons: <u>*Quantum excitations*</u> of solids

Measuring phonons

Inelastic neutron scattering
Inelastic light scattering (Raman/Brillouin)
(far) Infrared absorption
Electron energy loss spectroscopy
Inelastic X-ray scattering
Ultrasound propagation
Point and tunnel spectroscopy
…





Inelastic scattering



Energy conservation

$$\frac{\hbar^2 k^2}{2m_n} = \frac{\hbar^2 k'^2}{2m_n} \pm \hbar\Omega$$

"Momentum" conservation

$$k = k' \pm q + G$$

Diamond lattice





Phonon dispersion in Si





Si: FCC diamond structure (2 at. Prim. Cell)

Phonon dispersion in copper





Γ at (0,0,0)

Cu: FCC (1 at/cell)
Recipr. Space BCC

Phonons

- Quantized lattice vibrations
- #modes: 3s
- Optical & acoustical
- Transversal & longitudinal
- Relevant **k** vectors in first BZ
- Thermal occupation: Planck
- Dispersion $\omega(\mathbf{k})$
- Energy $\hbar\omega(\mathbf{k})$
- 'Momentum' $\hbar\mathbf{k}$
- Sound velocity

Thermal properties

Thermal properties

- Thermal expansion (classical)
- Lattice specific heat
  - Density of States
    - Debye model
    - Einstein model
- Thermal conductivity
  - Phonon scattering, mean free path

Thermal expansion

Harmonic crystal: No thermal expansion!!



$$r_j = ja + u_j$$

Harmonic

$$V(u_j) = \alpha \cdot u_j^2$$

$$a(T) = a + \langle u_j \rangle_T = a$$

$$\langle u_j \rangle_T = \frac{\int du \cdot u\, e^{-V(u_j)/k_B T}}{\int du \cdot e^{-V(u_j)/k_B T}} = 0$$



No lattice expansion !!

Anharmonicity: Thermal expansion (classical)



$$r_j = ja + u_j$$

Weak anharmonicity

$$V(u_j) = \alpha \cdot u_j^{\,2} - \gamma \cdot u_j^{\,3}$$

$$\langle u_j \rangle_T = \frac{\int du \cdot u\, e^{-V(u_j)/k_B T}}{\int du \cdot e^{-V(u_j)/k_B T}} \approx \frac{3/4\,(k_B T)^{3/2}\sqrt{\pi}\,\gamma\,\alpha^{-5/2}}{\sqrt{k_B T}\,\sqrt{\pi}\,\alpha^{-1/2}}$$

$$a(T) = a_0 + \langle u_j \rangle_T = a_0 + \frac{3\gamma}{4\alpha^2} k_B T$$



Lattice expansion is caused by anharmonicity !

Specific heat

Heat capacity: $C_v = \left.\frac{\partial U}{\partial T}\right|_V$

Summation over modes k and branches p:

$$U = \sum_{k,p} U_{k,p} = \sum_{k,p} \langle n_{k,p} \rangle \hbar\omega_{k,p} = \sum_{k,p} \frac{\hbar\omega_{k,p}}{e^{\hbar\omega_{k,p}/k_B T} - 1}$$

Number of modes in range $\omega \to \omega + d\omega$: $d\omega\, D_p(\omega)$

D(ω): Density of states

$$\sum_k \quad \to \quad \int d\omega\, D_p(\omega)$$

$$U = \sum_p \int d\omega D_p(\omega) \frac{\hbar\omega}{e^{\hbar\omega/k_B T} - 1}$$

Lattice heat capacity

$$\boxed{\sum_k \quad \rightarrow \quad \int d\omega \, D_p(\omega)}$$

$$\downarrow$$

$$U = \sum_p \sum_k U_{k,p} = \sum_p \sum_k \langle n_{k,p} \rangle \hbar\omega_{k,p} = \sum_p \int d\omega D_p(\omega) \frac{\hbar\omega}{e^{\hbar\omega/k_B T} - 1}$$

$$\rightarrow \quad \boxed{C_{lattice} = \frac{\partial U}{\partial T} = k_B \sum_p \int d\omega D_p(\omega) \frac{x^2 e^x}{(e^x - 1)^2}}$$

All lattice properties in D(ω):

<u>Density of states:</u> # modes per unit frequency

Density of states in 1D

1D crystal, N atoms, length L=Na

Vibrational mode: $u_k(j,t) = u_k e^{-i(\omega t - kaj)}$

Periodicity over L=Na: $u_k \cdot e^{ikaj} = u_k \cdot e^{ika(j+N)}$

$$\rightarrow \quad e^{ikaN} = 1 \qquad kaN = 2\pi s \qquad k = \frac{2\pi s}{Na} = \frac{2\pi}{L} s$$

$$-\frac{\pi}{a} < k \leq \frac{\pi}{a} \quad \Rightarrow \quad s = -\frac{N-1}{2}, \ldots, \frac{N-1}{2}, \frac{N}{2}$$

i.e. <u>N modes in the first BZ</u>

Density of states in 1D



modes/unit 'length' = N / (2π/a) = L/ 2π

In interval dk: $$D(k)dk = \frac{N}{2\pi/a}dk = \frac{L}{2\pi}dk$$

$$\boxed{D(\omega)d\omega = \frac{L}{2\pi}\frac{dk}{d\omega}d\omega}$$

Density of states (1D)



$$D(\omega)d\omega = \frac{L}{2\pi}\frac{dk}{d\omega}d\omega$$

1D, 1 at./cell: $\omega(k) = \sqrt{\frac{4C}{m}}\left|\sin(\frac{ka}{2})\right| \quad \Rightarrow \quad \omega(k) = \omega_m \left|\sin(ka/2)\right|$

$$\left[\frac{\partial\omega(k)}{\partial k}\right]^{-1} = \frac{2}{a}\frac{1}{\omega_m\left|\cos(ka/2)\right|} = \frac{2}{a}\frac{1}{\sqrt{\omega_m^2 - \omega^2}}$$

$$\Rightarrow \quad D(\omega) = 2\frac{N}{\pi}\frac{1}{\sqrt{1-\omega^2}}$$

Density of states (1D)



$$D(\omega) = 2\frac{N}{\pi}\frac{1}{\sqrt{\omega_m^2 - \omega^2}}$$





Density of states in 3D

3D crystal, N^3 atoms, cube length L

Periodic boundary conditions: $e^{i(k_x x + k_y y + k_z z)} = e^{i(k_x(x+L) + k_y(y+L) + k_z(z+L))}$

$$\rightarrow \quad k_x, k_y, k_z = s \cdot \frac{2\pi}{L}; \qquad s = -\frac{N-1}{2}, \ldots, \frac{N-1}{2}, \frac{N}{2}$$

One value of k per volume $\left(\frac{2\pi}{L}\right)^3$

Total # k values in sphere with radius k: $N(k) = \frac{V_k}{(2\pi/L)^3} = \frac{4\pi k^3/3}{(2\pi)^3/V}$

$$D(k) = \frac{dN(k)}{dk} = \frac{Vk^2}{2\pi^2} \qquad \boxed{D(\omega) = \frac{Vk^2}{2\pi^2}\frac{dk}{d\omega}}$$

Density of states in 3D



COPPER



Density of States

	$D(k)$	$D(\omega)$
1D	$\frac{L}{2\pi}$	$\frac{L}{2\pi}\frac{dk}{d\omega}$
2D	$\frac{A}{2\pi^2}k$	$\frac{A}{2\pi^2}k\frac{dk}{d\omega}$
3D	$\frac{V}{2\pi^2}k^2$	$\frac{V}{2\pi^2}k^2\frac{dk}{d\omega}$

Back to lattice heat capacity

$$C_{lattice} = \frac{\partial U}{\partial T} = k_B \sum_p \int d\omega D_p(\omega) \frac{x^2 e^x}{\left(e^x - 1\right)^2}$$

DOS in general complicated function
Simplifications:

- Debye model (take sound velocity constant)
- Einstein model (take phonon frequency constant)

Debije model for DOS

$$\omega = v \cdot k$$

Number of modes with |k'| < |k| for each polarization (branch):



$$N\left(|\vec{k}|\right) = \frac{\frac{4}{3}\pi|\vec{k}|^3}{\left(\frac{2\pi}{V_{cell}}\right)^3}$$

Number of modes: $N(k_D) = N_{atom}$



$$\Rightarrow k_D = \left(\frac{6\pi^2 N_{atom}}{V_{cell}}\right)^{1/3}$$

$$\Rightarrow N(\omega < \omega_D) = \frac{V_{cell}}{(2\pi)^3}\frac{4\pi}{3}\left(\frac{\omega}{v}\right)^3$$

Debije model

Debije Frequency: $$\omega_D = v\left(\frac{6\pi^2 N_{atom}}{V_{cell}}\right)^{1/3}$$

Debije Temperature: $$\theta = \frac{\hbar\omega_D}{k_B} = \frac{\hbar v}{k_B}\left(\frac{6\pi^2 N_{atom}}{V_{cell}}\right)^{1/3}$$

Density of states: $$D(\omega) = \frac{dN(\omega)}{d\omega} = V_{cell}\frac{\omega^2}{2\pi^2 v^3}$$

Total energy stored in phonons:

$$U = \sum_{k,p}\frac{\hbar\omega_{k,p}}{e^{\hbar\omega_{k,p}/k_BT}-1} = 3\int_0^{\omega_D}\frac{\hbar\omega}{e^{\hbar\omega/k_BT}-1}D(\omega)d\omega$$

Debije specific heat

Total energy :

$$U = \frac{3V_{cell}\hbar}{2\pi^2 v^3}\int_0^{\omega_D} \frac{\omega^3}{\exp(\hbar\omega / k_B T) - 1} d\omega$$

Specific heat:

$$C_{lat} = \frac{dU}{dT} = \frac{3V_{cell}\hbar}{2\pi^2 v^3}\int_0^{\omega_D} \frac{\omega^3 \exp(\hbar\omega / k_B T)}{\left(\exp(\hbar\omega / k_B T) - 1\right)^2} \frac{\hbar\omega}{k_B T^2} d\omega$$

$$C_{lat} = \frac{dU}{dT} = 9N_{atom}\left(\frac{T}{\theta}\right)^3 \cdot \int_0^{\theta/T} \frac{x^4 e^x}{\left(e^x - 1\right)^2} dx \qquad x = \frac{\hbar\omega}{k_B T}$$

Debije T^3 law

Low temperature limit: $\theta / T \to \infty$

$$\int_0^\infty \frac{x^4 e^x}{(e^x - 1)^2} dx = \frac{\pi^4}{15}$$

$$\boxed{C_{lat} = \frac{12\pi^4}{5} N_{atom} k_B \left(\frac{T}{\theta}\right)^3}$$

High temperature limit: $\theta / T \to 0$

$$\int_0^{\theta/T} \frac{x^4 e^x}{(e^x - 1)^2} dx \approx \frac{1}{3}\left(\frac{\theta}{T}\right)^3$$

$$\boxed{C_{lat} = 3 N_{atom} k_B}$$ (Dulong-Petit, $U=3N \cdot k_B T$)

Specific heat



Specific heat AlN



Specific heat Aluminum Nitride (Koshchenko et al., 1985)

Specific heat Cu

Copper again





Specific heat $Th_xU_{1-x}Be_{13}$



Heavy fermion system, 2 S.C. transitions (Z. Fisk et al., 2000)